

AD-A153 548

ON APPROXIMATING A SET-VALUED FUNCTION LOCALLY(U)  
WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER  
V F DEMYANOV ET AL. JAN 85 MRC-TSR-2786

1/1


UNCLASSIFIED

DAAG29-88-C-0041

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS 1963-A

MRC Technical Summary Report #2/86

ON APPROXIMATING A SET-VALUED FUNCTION  
LOCALLY

V. F. Demyanov, C. Lemarechal and  
J. Zowe

**Mathematics Research Center  
University of Wisconsin—Madison  
610 Walnut Street  
Madison, Wisconsin 53705**

January 1985

(Received December 19, 1984)

**Approved for public release  
Distribution unlimited**

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

DTIC  
ELECT  
MAY 9 1985  
S D

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

ON APPROXIMATING A SET-VALUED FUNCTION LOCALLY

V. F. Demyanov<sup>1</sup>, C. Lemarechal<sup>2</sup> and J. Zowe<sup>3</sup>

Technical Summary Report #2786  
January 1985

ABSTRACT

Given a multi-valued mapping  $F$ , we address the problem of finding another multi-valued mapping  $H$  that agrees locally with  $F$  in some sense. We show that, contrary to the scalar case, introducing a derivative of  $F$  is hardly convenient. For the case when  $F$  is convex-compact-valued, we give some possible approximations, and at the same time we show their limitations. The present paper is limited to informal demonstration of concepts and mechanisms. Formal statements and their proofs will be published elsewhere.

AMS (MOS) Subject Classifications: 41-01, 46G05

Key Words: convexity, multifunctions, differentiability, Newton scheme.

Work Unit Number 5 (Optimization and Large Scale Systems)

<sup>1</sup>IIASA, 2361 Laxenburg (Austria)

<sup>2</sup>INRIA, B.P. 105, 78153 Le Chesnay (France)

<sup>3</sup>University of Bayreuth, P.F. 3008, 8580 Bayreuth (W. Germany)

Part of this research was performed at the Mathematics Research Center of the University of Wisconsin-Madison, and sponsored by the United States Army under Contract No. DAFG29-80-C-0041.

Distribution For	
NTIS GRA&I <input checked="" type="checkbox"/>	
DTIC TAB <input type="checkbox"/>	
Unannounced <input type="checkbox"/>	
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## SIGNIFICANCE AND EXPLANATION

This paper is concerned with extension of the concept of derivative from functions to multi-valued mappings. Proper definitions of such extensions are useful to solve inclusions, and more specifically to minimize convex functions. Simple examples are given to show the difficulties, and some proposals are made to overcome them.

---

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

# ON APPROXIMATING A SET-VALUED FUNCTION LOCALLY

V. F. Demyanov<sup>1</sup>, C. Lemarechal<sup>2</sup> and J. Zowe<sup>3</sup>

## 1. INTRODUCTION

Consider first the problem of solving a nonlinear system:

$$f(x) = 0 \quad (1)$$

where  $f$  is a vector-valued function. If we find a *first order approximation* of  $f$  near  $x$ , i. e. a vector-valued bi-function  $h$  such that

$$h(x;d) = f(x+d) + o(d) \quad (2)$$

(where  $o(d)/\|d\| \rightarrow 0$  when  $d \rightarrow 0$ ) then we can apply the *Newton principle*: given a current iterate  $x$ , solve for  $d$

$$h(x;d) = 0 \quad (3)$$

(supposedly simpler than (1)) and move to  $x+d$ .

Everybody knows that if  $f$  is differentiable and if, in addition to satisfying (2),  $h$  is required to be affine in  $d$ , then it is *unambiguously* defined by

$$h(x;d) := f(x) + f'(x)d \quad (4)$$

Merging (2) and (4) and subtracting  $f(x)$  gives also a nonambiguous definition of  $f'$  (the jacobian operator of  $f$ ) by:

$$f'(x)d := f(x+d) - f(x) + o(d).$$

Suppose now that we have to solve

$$0 \in F(x) \quad (5)$$

where  $F$  is a multi-valued mapping, i. e.  $F(x) \subset \mathbb{R}^n$ . A possible application of (5) is in nonsmooth optimization, when  $F$  is the (approximate) subdifferential of an objective function to be minimized. To apply the same principle as in the single valued case,  $F(x+d)$  must be approximated by some set

---

<sup>1</sup>IIASA, 2361 Laxenburg (Austria)

<sup>2</sup>INRIA, B.P. 105, 78153 Le Chesnay (France)

<sup>3</sup>University of Bayreuth, P.F. 3008, 8580 Bayreuth (W.Germany)

$H(x;d) \subset \mathbb{R}^n$ . Continuing the parallel and requiring  $H$  to be affine in  $d$  (whatever it means), we must express it as a *sum of two sets*:  $H(x,d) = F(x) + G$ . In summary, we want to find a set  $G$  such that, for all  $\varepsilon > 0$  and  $\|d\|$  small enough:

$$F(x+d) \subset F(x) + G + \varepsilon \|d\| U \quad (6.a)$$

and

$$F(x) + G \subset F(x+d) + \varepsilon \|d\| U \quad (6.b)$$

where  $U$  is the unit ball of  $\mathbb{R}^n$ . Unfortunately, such a writing is already worthless. First, it does not help defining the "linearization"  $G$ : just because the set of subsets is not a group,  $F(x)$  cannot be subtracted in (6). Furthermore, (6) is extremely restrictive: for  $n = 1$ , consider the innocent mapping  $F(x) := [0, 3x]$  (defined for  $x \geq 0$ ). Take  $x = 1$ ,  $\varepsilon = 1$  and  $d < 0$ . It is impossible to find a set  $G$  satisfying (6.b), i. e.  $[0, 3] + G \subset [d, 3+2d]$ . For example,  $G = \{d\}$  is already too "thick".

A conclusion of this section is that a first order approximation to a multivalued mapping cannot be readily constructed by a standard linearization; the definition of such an approximation is at present ambiguous. For a deep insight into differentiability of sets, we refer to [6] and its large bibliography. Here, for want of a complete theory, we will give in the next sections two possible proposals. None of them is fully satisfactory, but they are rather complementary, in the sense that each one has a chance to be convenient when the other is not. We will restrict ourselves to the convex compact case. Furthermore, as is usual in nondifferentiable optimization, we will consider only directional derivatives. Therefore we adopt simpler notations:  $x$  and the direction  $d$  being fixed, we call  $F(t)$  the image by  $F$  of  $x + td$ ,  $t \geq 0$ . We say that  $H$  approximates  $F$  to 1st order near  $t = 0^+$  if for every  $\varepsilon > 0$ , there is  $\delta > 0$  such that  $t \in [0, \delta]$  implies

$$F(t) \subset H(t) + \varepsilon t U \text{ and } H(t) \subset F(t) + \varepsilon t U \quad (7)$$

Note that, among others,  $F$  approximates itself!

## 2. MAPPINGS DEFINED BY A SET OF CONSTRAINTS

As a first illustration, suppose  $F$  is defined by:

$$F(t) := \{z \in \mathbb{R}^n \mid c_j(t, z) \leq 0 \text{ for } j = 1, \dots, m\}$$

where the "constraints"  $c_j$  are convex in  $z$ . Assume the existence of  $c'_j(0, z)$ , the right derivative of  $c_j(\cdot, z)$  at  $t = 0$  ( $c'_j(0^+, z)$  would be more suggestive). Then it is natural to consider approximating  $F(t)$  by

$$H(t) := \{z \mid c_j(0, z) + t c'_j(0, z) \leq 0 \text{ for } j = 1, \dots, m\}. \quad (8)$$

An algorithm based on this set would then be quite in the spirit of [7].

It is possible to prove that the  $H$  of (8) does satisfy (7), provided some hypotheses hold, for example

- (i)  $[c_j(t, z) - c_j(0, z)] / t \rightarrow c'_j(0, z)$  uniformly in  $z$ , when  $t \downarrow 0$ ,
- (ii) there exists  $z_0$  such that  $c_j(0, z_0) < 0$  for  $j = 1, \dots, m$ .

A weak point of (8) is that it is highly non-canonical. For example, perturbing the constraints to  $(1 + a_j t) c_j(t, z)$  gives the same  $F$  but does change  $H$ .

### 3. A DIRECT SET-THEORETIC CONSTRUCTION

If we examine (6) again, we see that there would be no difficulty if  $F(x)$  were a singleton: then (6) would always be consistent because  $F(x+d)$  would never be less thick than  $F(x)$ , and  $F(x)$  could be subtracted. This leads to differentiating  $F$  at an arbitrary but fixed  $y \in F(0)$ . Define

$$F'_y(0) := \left\{ z \mid \begin{array}{l} \text{there exist } t_n \text{ and } y_n \in F(t_n) \text{ for } n \in \mathbb{N} \\ \text{with } t_n \downarrow 0 \text{ and } (y_n - y) / t_n \rightarrow z \end{array} \right\}$$

or, in a set-theoretic notation (see [2], Chapter VI):

$$F'_y(0) := \limsup_{t \downarrow 0} [F(t) - y] / t$$

This set is called the *contingent derivative* in [1], the (radial) *upper Dini derivative* in [6] and the *feasible set of first order* in [3]. We refer to [1] for an extensive study of  $F'$ , but some remarks will be useful:

- a)  $F'_y(0)$  depends on the behaviour of  $F$  near  $y$  only. If we take an arbitrary  $\alpha > 0$  and set  $G(t) := F(t) \cap \{y + \alpha U\}$ , then  $G'_y(0) = F'_y(0)$ .
- b) If  $F(t) \equiv F(0)$  does not depend on  $t$ ,  $F'_y(0)$  is just the tangent cone to  $F(0)$  at  $y$ .
- c) Let  $A$  be a convex set in  $\mathbb{R}^n$ , and  $f: [0, 1] \rightarrow \mathbb{R}^n$  a differentiable mapping (with  $f(0) = 0$  for notational simplicity). Consider  $F(t) := \{f(t)\} + A$ . Given  $y \in F(0) = A$ , call  $T_y$  the tangent cone to  $F(0) = A$  at  $y$ . Then it can be shown that  $F'_y(0) = \{f'(0)\} + T_y$ . This is the situation when  $F$  is the approximate subdifferential of a convex quadratic function (see [4]).
- d) Let  $n = 2$ . Given  $r \in \mathbb{R}$ , consider  $F(t) := P(t) \cap U$  with the halfspace  $P(t) := \{y = (y_1, y_2) \mid y_2 \geq r t y_1\}$ . It can be shown that, for  $y = 0 \in F(0)$ ,  $F'_0(0) = \{z = (z_1, z_2) \mid z_2 \geq 0\}$ ;  $F'_0(0)$  is the same as it would be if  $r$  were 0 (in which case  $F(t)$  would be fixed), and does not predict the rotation of  $F(t)$  around  $y = 0$ .



Because a convex set is the intersection of the cones tangent to it, our remark b) above suggests to approximate  $F(t)$  by

$$H(t) := \cap \{y + tF'_0(O) \mid y \in F(O)\} \quad (9)$$

Of course, this will be possible only under additional assumptions (not only due to the multi-valuedness of  $F$ ; for example  $F(t) := \{t \sin \log t\}$  has  $F(O) = \{O\}$ ,  $F'_0(O) = [-1, +1]$  and  $H(t) = [-t, +t]$ ).

Before mentioning the assumptions in question, we introduce another candidate to approximate  $F$ : for  $p \in \mathbb{R}^n$ , denote by  $s_p(t) := \sup \{ \langle p, y \rangle \mid y \in F(t) \}$  the support function of  $F(t)$ . It is known that  $F$  can be described in terms of  $s$ , namely  $F(t) = \{y \mid \langle p, y \rangle \leq s_p(t) \ \forall p \in \mathbb{R}^n\}$ . Then, if  $s_p$  has a (directional) derivative  $s'_p(O)$ , the following set is natural (see [5]):

$$G(t) := \{y \mid \langle p, y \rangle \leq s_p(O) + t s'_p(O) \ \forall p \in \mathbb{R}^n\}. \quad (10)$$

To assess these candidates (9) and (10), the following assumptions can be considered:

- (i)  $[s_p(t) - s_p(O)]/t \rightarrow s'_p(O)$  uniformly for  $p \in U$ , when  $t \downarrow 0$ ;
- (ii)  $F(O)$  has a nonempty interior.

They allow to prove:

If (i) holds, then  $H(t) = G(t)$ ; if (ii) also holds, then (7) holds.

We remark that (i) alone suffices to prove the second half of (7), which is the important one for (5) (solving  $O \in H(t)$  gives some among the possible Newton iterates); however  $H(t)$  may be void if (ii) does not hold. It is also interesting to remark that, if  $s'_p(O)$  is assumed to be convex in  $p$  (in which case (ii) is not needed), then it is the support function of a convex set that we are entitled to call  $F'(O)$  because there holds  $H(t) = F(O) + t F'(O)$  (due to additivity of support functions). In other words, convexity of  $s'_p(O)$  gives the "easy" situation in which (6) holds.

The role of assumption (i) is more profound. It is natural to require that  $F'_y(O)$  does predict the behaviour of  $F(t)$  near  $y$ ; this behaviour is trivial when  $y \in \text{int } F(O)$  (then  $F(t)$  must contain  $y$  for all  $t$  small enough); if  $y$  is on the boundary of  $F(O)$  then there is a normal cone  $N_y(O)$  to  $F(O)$  at  $y$ , and  $s_p(O) = \langle p, y \rangle$  for  $p \in N_y(O)$ ; hence the behaviour of  $F(t)$  near  $y$  is naturally related to the behaviour of  $s_p(t)$  for these normal  $p$ 's (incidentally, a key result is that  $F'_y(O) = \{z \mid \langle p, z \rangle \leq s'_p(O) \ \forall p \in N_y(O)\}$ ; (i) is essential for this). However, it is not only some technicalities in the proof that require the *uniformity* stated in (i), but rather the deficiency of  $F'$  suggested by d) above: consider the innocent mapping


$$F(t) := \{y = (y_1, y_2) \mid 0 \leq y_1 \leq 1, ty_1 \leq y_2 \leq 1\}.$$

Given  $\alpha \in \mathbb{R}$  and  $p = (\alpha, -1)$ ,  $s_p(t) = \max \{(\alpha - t)y_1 \mid 0 \leq y_1 \leq 1\}$  and thus, (i) is violated: when  $\alpha \downarrow 0$ ,  $s'_p(0)$  jumps from  $-1$  to  $0$ . For this example,  $H(t) = G(t) = [0, 1] \times [t, 1]$ , which is a poor approximation of  $F(t)$ . This is rather disappointing, but observe that Section 2 is well-suited for the present  $F$ .

# REFERENCES

- [1] J. P. Aubin, "Contingent derivatives of set-valued maps and existence of solutions to nonlinear inclusions and differential inclusions". *MRC Technical Summary Report 2044*, University of Wisconsin, Madison (1980). See also: same title, in: L. Nachbin (ed.) *Mathematical Analysis and Applications*, Academic Press (1981) 159 - 229.
- [2] C. Berge, *Topological Spaces*, Mac Millan, London, 1963.
- [3] V. F. Demyanov and I. M. Lupikov, "Extremal functions over the  $\epsilon$ -sub-differential mapping", *Vestnik Leningrad University* 1 (1983) 27 - 32.
- [4] J. B. Hiriart-Urruty, " $\epsilon$ -subdifferential calculus", in: J. P. Aubin, R. B. Vinter (eds.) *Convex Analysis and Optimization*, Pitman (1982) 43 - 92.
- [5] C. Lemaréchal and E. A. Nurminskii, "Sur la différentiabilité de la fonction d'appui du sous-différentiel approché", *C. R. Acad. Sci. Paris* 290, 18 (1980) 855 - 858.
- [6] J. P. Penot, "Differentiability of relations and differential stability of perturbed optimization problems", *SIAM Control* 22, 4 (1984) 529 - 551.
- [7] S. M. Robinson, "Extension of Newton's method to nonlinear functions with values in a cone", *Numerische Mathematik* 19 (1972) 341 - 347.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2786	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) ON APPROXIMATING A SET-VALUED FUNCTION LOCALLY		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) V. F. Dem'yanov, C. Lemarechal and J. Zowe		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 5 - Optimization and Large Scale Systems
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE January 1985
		13. NUMBER OF PAGES 6
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) convexity, multifunctions, differentiability, Newton scheme		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Given a multi-valued mapping $F$ , we address the problem of finding another multi-valued mapping $H$ that agrees locally with $F$ in some sense. We show that, contrary to the scalar case, introducing a derivative of $F$ is hardly convenient. For the case when $F$ is convex-compact-valued, we give some possible approximations, and at the same time we show their limitations. The present paper is limited to informal demonstration of concepts and mechanisms. Formal statements and their proofs will be published elsewhere.		



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A